PROCEEDINGS OF SCIENCE

Searches for Lepton Number Violation and resonances in the $K^{\pm} \to \pi\mu\mu$ decays at the NA48/2 experiment

Karim Massri[*†]
University of Liverpool
E-mail: karim.massri@cern.ch

The NA48/2 experiment at CERN collected a large sample of charged kaon decays into final states with multiple charged particles in 2003–2004. A new upper limit on the rate of the lepton number violating decay $K^{\pm} \to \pi^{\mp}\mu^{\pm}\mu^{\pm}$ obtained from this sample is reported: $\mathscr{B}(K^{\pm} \to \pi^{\mp}\mu^{\pm}\mu^{\pm}) < 8.6 \times 10^{-11}$ at 90% CL. Searches for two-body resonances in the $K^{\pm} \to \pi\mu\mu$ decays (including heavy neutral leptons N_4 and inflatons χ) in the accessible range of masses and lifetimes are also presented. In the absence of a signal, upper limits are set on the products of branching ratios $\mathscr{B}(K^{\pm} \to \mu^{\pm}N_4)\mathscr{B}(N_4 \to \pi\mu)$ and $\mathscr{B}(K^{\pm} \to \pi^{\pm}\chi)\mathscr{B}(\chi \to \mu^{+}\mu^{-})$ as functions of the resonance mass and lifetime. These limits are in the $10^{-10} - 10^{-9}$ range for resonance lifetimes below 100 ps.

38th International Conference on High Energy Physics
3-10 August 2016
Chicago, USA

[*]Speaker.

[†]On behalf of the NA48/2 Collaboration: G. Anzivino, R. Arcidiacono, W. Baldini, S. Balev, J.R. Batley, M. Behler, S. Bifani, C. Biino, A. Bizzeti, B. Bloch-Devaux, G. Bocquet, N. Cabibbo, M. Calvetti, N. Cartiglia, A. Ceccucci, P. Cenci, C. Cerri, C. Cheshkov, J.B. Chèze, M. Clemencic, G. Collazuol, F. Costantini, A. Cotta Ramusino, D. Coward, D. Cundy, A. Dabrowski, P. Dalpiaz, C. Damiani, M. De Beer, J. Derré, H. Dibon, L. DiLella, N. Doble, K. Eppard, V. Falaleev, R. Fantechi, M. Fidecaro, L. Fiorini, M. Fiorini, T. Fonseca Martin, P.L. Frabetti, L. Gatignon, E. Gersabeck, A. Gianoli, S. Giudici, A. Gonidec, E. Goudzovski, S. Goy Lopez, M. Holder, P. Hristov, E. Iacopini, E. Imbergamo, M. Jeitler, G. Kalmus, V. Kekelidze, K. Kleinknecht, V. Kozhuharov, W. Kubischta, G. Lamanna, C. Lazzeroni, M. Lenti, L. Litov, D. Madigozhin, A. Maier, I. Mannelli, F. Marchetto, G. Marel, M. Markytan, P. Marouelli, M. Martini, L. Masetti, E. Mazzucato, A. Michetti, I. Mikulec, N. Molokanova, E. Monnier, U. Moosbrugger, C. Morales Morales, D.J. Munday, A. Nappi, G. Neuhofer, A. Norton, M. Patel, M. Pepe, A. Peters, F. Petrucci, M.C. Petrucci, B. Peyaud, M. Piccini, G. Pierazzini, I. Polenkevich, Yu. Potrebenikov, M. Raggi, B. Renk, P. Rubin, G. Ruggiero, M. Savrié, M. Scarpa, M. Shieh, M.W. Slater, M. Sozzi, S. Stoynev, E. Swallow, M. Szleper, M. Valdata-Nappi, B. Vallage, M. Velasco, M. Veltri, S. Venditti, M. Wache, H. Wahl, A. Walker, R. Wanke, L. Widhalm, A. Winhart, R. Winston, M.D. Wood, S.A. Wotton, A. Zinchenko, M. Ziolkowski.

© Copyright owned by the author(s) under the terms of the Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International License (CC BY-NC-ND 4.0).

arXiv:1611.07375v1 [hep-ex] 22 Nov 2016

1. Introduction

The NA48/2 experiment at CERN SPS was a multi-purpose $K^{\pm}$ experiment which collected data in 2003–2004, whose main goal was to search for direct CP violation in the $K^{\pm} \to \pi^{\pm}\pi^{+}\pi^{-}$ and $K^{\pm} \to \pi^{\pm}\pi^{0}\pi^{0}$ decays [1]. The large statistics of the samples of charged kaon decays into final states with multiple charged particles collected allows to search for the forbidden LNV $K^{\pm} \to \pi^{\mp}\mu^{\pm}\mu^{\pm}$ decay, as well as for two-body resonances in $K^{\pm} \to \pi\mu\mu$ decays. Since a particle X produced in a $K^{\pm} \to \mu^{\pm}X$ ($K^{\pm} \to \pi^{\pm}X$) decay and decaying promptly to $\pi^{\pm}\mu^{\mp}$ ($\mu^{+}\mu^{-}$) would produce a narrow spike in the invariant mass $M_{\pi\mu}$ ($M_{\mu\mu}$) spectrum, the invariant mass distributions of the collected $K^{\pm} \to \pi\mu\mu$ samples have been scanned looking for such a signature.

2. Selected data samples

The event selection is based on the reconstruction of a three-track vertex: given the resolution of the vertex longitudinal position ($\sigma_{vtx} = 50$ cm), $K^{\pm} \to \pi^{\mp}\mu^{\pm}\mu^{\pm}$ and $K^{\pm} \to \pi^{\pm}\mu^{+}\mu^{-}$ decays (denoted $K^{\rm LNV}_{\pi\mu\mu}$ and $K^{\rm LNC}_{\pi\mu\mu}$ below) mediated by a short-lived ($\tau \lesssim 10$ ps) resonant particle are indistinguishable from a genuine three-track decay. The size of the selected $K_{\pi\mu\mu}$ samples is normalised relative to the abundant $K^{\pm} \to \pi^{\pm}\pi^{+}\pi^{-}$ channel (denoted $K_{3\pi}$ below), from which the number of $K^{\pm}$ decays in the 98 m long fiducial decay region is obtained: $N_K = (1.64 \pm 0.01) \times 10^{11}$. The $K_{\pi\mu\mu}$ and $K_{3\pi}$ samples are collected concurrently using the same trigger logic.

The invariant mass distributions of data and MC events passing the $K^{\rm LNV}_{\pi\mu\mu}$ and $K^{\rm LNC}_{\pi\mu\mu}$ selections are shown in Fig. 1. One event is observed in the signal region after applying the $K^{\rm LNV}_{\pi\mu\mu}$



Figure 1: Invariant mass distributions of data and MC events passing the $K^{\rm LNV}_{\pi\mu\mu}$ (left) and $K^{\rm LNC}_{\pi\mu\mu}$ (right) selections. The signal mass regions are indicated with vertical arrows.

selection, while 3489 $K^{\rm LNC}_{\pi\mu\mu}$ candidates are selected with the $K^{\rm LNC}_{\pi\mu\mu}$ selection. A peak search assuming different mass hypotheses is performed over the distributions of the invariant masses M_{ij} ($ij = \pi^{\pm}\mu^{\mp}, \mu^{+}\mu^{-}$) of the selected $K_{\pi\mu\mu}$ samples. In total, 284 (267) and 280 mass hypotheses

are tested respectively for the search of resonances in the $M_{\pi\mu}$ distribution of the $K^{\rm LNV}_{\pi\mu\mu}$ ($K^{\rm LNC}_{\pi\mu\mu}$) candidates and in the $M_{\mu\mu}$ distribution of the $K^{\rm LNC}_{\pi\mu\mu}$ candidates, covering the full kinematic ranges.

3. Results

3.1 Upper Limit on $\mathscr{B}(K^{\pm} \to \pi^{\mp}\mu^{\pm}\mu^{\pm})$

The upper limit (UL) at 90% confidence level (CL) on the number of $K^{\pm} \to \pi^{\mp}\mu^{\pm}\mu^{\pm}$ signal events in the $K^{\rm LNV}_{\pi\mu\mu}$ sample corresponding to the observation of one data event and a total number of expected background events $N_{bkg} = 1.163 \pm 0.867_{stat} \pm 0.021_{ext} \pm 0.116_{syst}$ is obtained applying an extension of the Rolke-Lopez method [2]: $N^{\rm LNV}_{\pi\mu\mu} < 2.92$ at 90% CL. Using the values of the signal acceptance $A(K^{\rm LNV}_{\pi\mu\mu}) = 20.62\%$ estimated with MC simulations and the number N_K of kaon decays in the fiducial volume (Sec. 2), the UL on the number of $K^{\pm} \to \pi^{\mp}\mu^{\pm}\mu^{\pm}$ signal events in the $K^{\rm LNV}_{\pi\mu\mu}$ sample leads to a constraint on the signal branching ratio $\mathscr{B}(K^{\pm} \to \pi^{\mp}\mu^{\pm}\mu^{\pm})$:

$$\mathscr{B}(K^{\pm} \to \pi^{\mp}\mu^{\pm}\mu^{\pm}) = \frac{N^{\rm LNV}_{\pi\mu\mu}}{N_K \cdot A(K^{\rm LNV}_{\pi\mu\mu})} < 8.6 \times 10^{-11} \quad @\ 90\%\ \text{CL}. \tag{3.1}$$

3.2 Results of the search for two-body resonances

No signal is observed, as the local significances of the signals in each mass hypothesis never exceed 3 standard deviations. In absence of a signal, ULs on the product $\mathscr{B}(K^{\pm} \to p_1 X)\mathscr{B}(X \to p_2 p_3)$ ($p_1p_2p_3 = \mu^{\pm}\pi^{\mp}\mu^{\pm}, \mu^{\pm}\pi^{\pm}\mu^{\mp}, \pi^{\pm}\mu^{+}\mu^{-}$) as a function of the resonance lifetime τ are obtained for each mass hypothesis m_i, by using the values of the acceptances $A_{\pi\mu\mu}(m_i, \tau)$ and the ULs on the number N^i_{sig} of signal events for such a mass hypothesis:

$$\mathscr{B}(K^{\pm} \to p_1 X)\mathscr{B}(X \to p_2 p_3)\big|_{m_i,\tau} = \frac{N^i_{sig}}{N_K \cdot A_{\pi\mu\mu}(m_i, \tau)}. \tag{3.2}$$

The obtained ULs on $\mathscr{B}(K^{\pm} \to p_1 X)\mathscr{B}(X \to p_2 p_3)$ ($p_1p_2p_3 = \mu^{\pm}\pi^{\mp}\mu^{\pm}, \mu^{\pm}\pi^{\pm}\mu^{\mp}, \pi^{\pm}\mu^{+}\mu^{-}$) as a function of the resonance mass, for several values of the resonance lifetime, are shown in Fig. 2.

4. Conclusions

The searches for the LNV $K^{\pm} \to \pi^{\mp}\mu^{\pm}\mu^{\pm}$ decay and resonances in $K^{\pm} \to \pi\mu\mu$ decays at the NA48/2 experiment, using the 2003–2004 data, are presented. No signals are observed. An UL of 8.6×10^{-11} for $\mathscr{B}(K^{\pm} \to \pi^{\mp}\mu^{\pm}\mu^{\pm})$ has been established, which improves the best previous limit [3] by more than one order of magnitude. ULs are set on the products $\mathscr{B}(K^{\pm} \to \mu^{\pm}N_4)\mathscr{B}(N_4 \to \pi\mu)$ and $\mathscr{B}(K^{\pm} \to \pi^{\pm}\chi)\mathscr{B}(\chi \to \mu^{+}\mu^{-})$ as functions of the resonance mass and lifetime. These limits are in the $10^{-10} - 10^{-9}$ range for resonance lifetimes below 100 ps.

References

[1] J.R. Batley *et al.* (NA48/2 collaboration), Eur. Phys. J. **C52** (2007) 875.

[2] W.A. Rolke and A.M. López, Nucl. Instrum. Meth. **A458** (2001) 745.

[3] J.R. Batley *et al.* (NA48/2 collaboration), Phys. Lett. **B697** (2011) 107.


a
b
c
τ = 100 ns
τ = 10 ns
τ = 1 ns
τ = 100 ps
τ < 10 ps
Assumed N4 mass, MeV/c2
Assumed X mass, MeV/c2
UL on BR(K± → μ±N4) × BR(N4 → π∓μ±) at 90% CL
UL on BR(K± → π±X) × BR(X → μ+μ−) at 90% CL

Figure 2: Obtained ULs at 90% CL on the products of branching ratios as functions of the resonance mass and lifetime: a) $\mathscr{B}(K^{\pm} \to \mu^{\pm} N_4)\mathscr{B}(N_4 \to \pi^{\mp}\mu^{\pm})$; b) $\mathscr{B}(K^{\pm} \to \mu^{\pm} N_4)\mathscr{B}(N_4 \to \pi^{\pm}\mu^{\mp})$; c) $\mathscr{B}(K^{\pm} \to \pi^{\pm}\chi)\mathscr{B}(\chi \to \mu^{+}\mu^{-})$. All presented quantities are strongly correlated for neighbouring resonance masses as the mass step of the scan is about 8 times smaller than the signal window width.